



## MD Accelerator

29.09.2016

<b>Liquid</b>	
Color	transparent
Smell	typical
Density	0,84 g/cm <sup>3</sup>
Viscosity at 20°C	1,05 mPa.s
Boiling temperature	78 °C
Flash point	12 °C
Vapour pressure	57 mbar
Water hazard class	2
Temperature resistance	+5°C to +40°C
<p>Marston-Domsel's accelerator ensures even faster hardening of MD GLUE instant adhesive. Mainly for use with instant adhesives with higher viscosities.</p>	
<b>Application</b>	
<p>Remove contamination and oil/grease from surfaces to be bonded using a product such as Marston Cleaner. Apply the MD activator for CA by spraying it on to one surface to be bonded. Apply adhesive to the other side and press the parts together immediately.</p>	
<b>Packaging</b>	<b>Item number</b>
25 glass bottles with spray head á 15 ml	MAC.Y15
<p>The values are average values. They serve merely for your information, but assume no warranty.</p>	

Bergheimer Str. 15 | D-53909 Zülpich | Tel. 02252/94150 | [info@marston-domsel.de](mailto:info@marston-domsel.de)  
[www.marston-domsel.de](http://www.marston-domsel.de)

The information in this product has been compiled to the best of our knowledge and is intended purely for information purposes. No claims can be inferred therefrom. Before use, thorough experiments should be carried out. Our brochure represents a basis. Responsibility for possible measures to protect property and persons lies with the user. Safety data sheets on the required standard are available for all products on request.